

EAGLE® 0.39% Granular Turf Fungicide

A granular product containing a systemic protectant and curative fungicide for the control of diseases in established turfgrass.

ACTIVE INGREDIENT:

myclobutanil: a-butyl-a-(4-chlorophenyl)-1-H-1,2,4-triazole-propanenitrile.....0.39%

INERT INGREDIENTS:..... 99.61%

Total.....100.00%

This product contains 0.156 lb of the active ingredient myclobutanil per 40 lb bag.

KEEP OUT OF REACH OF CHILDREN CAUTION - PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If in eyes	<ul style="list-style-type: none"> Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

NOTICE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-922-5994 day or night, for emergency treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

User must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or disposing of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

GENERAL INFORMATION

LESCO® Eagle® 0.39% Granular Turf Fungicide is a systemic, protectant and curative fungicide recommended for the control of listed diseases in established turfgrass. Examples of such sites include, but are not limited to: residential and commercial lawns, ornamental turf, grounds or lawns around business and office complexes, golf courses (fairways, roughs, tees, greens) and sod farms. Optimum disease control is achieved when this product is applied to established turfgrasses in a regularly scheduled preventative program.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made from any waterproof material
- Shoes plus socks.

NON-AGRICULTURAL USES REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Seed treatments and professional applications to lawn grasses, golf courses, industrial (office park), municipal and residential lawns are not within the scope of the Worker Protection Standard.

Keep unprotected persons out of treated area until dusts have settled.

USE DIRECTIONS FOR TURFGRASS

- LESCO® Eagle® 0.39% Granular Turf Fungicide should be used in conjunction with turf management practices that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing the disease. Diagnostic kits, extension experts or other identification methods should be used when developing disease control strategies.
- In non-residential turfgrass, optimum disease control is achieved when LESCO® Eagle® 0.39% Granular Turf Fungicide is applied in a preventative disease control program at a rate of 4 to 8 lb per 1000 sq ft. See the following table for specific application rates for various diseases.
- In residential turfgrass, optimum disease control is achieved when LESCO® Eagle® 0.39% Granular Turf Fungicide is applied in a preventative disease control program at a rate of 4 lb per 1000 sq ft. See the following table for specific application rates for various diseases.
- For the control of root and crown rot, this product must be watered-in using a minimum of 3 to 4 gallons of water per 1000 sq ft.
- Under conditions that are favorable for disease development, the interval between applications of LESCO® Eagle® 0.39% Granular Turf Fungicide should be reduced.
- Unless otherwise specified, when disease pressure is high of when used as a curative use higher rates of LESCO® Eagle® 0.39% Granular Turf Fungicide and shorter application interval. Under light to moderate disease pressure, apply LESCO® Eagle® 0.39% Granular Turf Fungicide at the low use rate and/or longer application intervals.
- Apply LESCO® Eagle® 0.39% Granular Turf Fungicide uniformly using an appropriate spreader. Calibrate the spreader before application as calibration may change with spreader age and use, walking speed and evenness of terrain. To calibrate the spreader, weigh out the amount of product necessary to treat an area of 800 sq ft. Using the suggested spreader setting provided by the manufacturer or from the table below, apply to a pre-measured area of 500 sq ft (5 x 100 sq ft) area. Weigh back product remaining in application equipment and adjust the spreader as necessary to give the correct rate of delivery for the measured area.
- To avoid pick-up, lightly irrigate treated areas soon after application.
- On short cut bentgrass (1/2" or less) when temperatures are above 80°F, apply only to dry foliage.

SUGGESTED SPREADER SETTINGS**

Recommended Rates (LESCO Eagle 0.39% Granular Turf Fungicide) in lb per 1,000 sq ft

Spreader	4 lb	6 lb	8 lb
LESCO	13	15	17
Cyclone® or Spiker®	3	4 1/2	4 1/2
Vibro®	20	28	32
Lely®	2 1/2 II	4 1/2 II	4 1/2 II

Apply LESCO® Eagle® 0.39% Granular Turf Fungicide with drop or rotary spreaders designed to apply granules. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions such that the spreader delivers the recommended application rate. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area in a given direction and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

#092019

T

EAGLE® 0.39% Granular Turf Fungicide

Non-Residential Turfgrass

Disease	LESCO® Eagle® 0.39% Granular Turf Fungicide (8/1000 sq ft)	Application Interval	Instructions	Restrictions
Anthracnose Red Thread Septoria Leaf Spot Brown Patch	4	14-21 days	Apply when conditions are favorable for disease development.	Do not apply more than 46 lb product per 1000 sq ft per year.
Dollar Spot Zona Leaf Spot	4	14 days	Begin applications when conditions are favorable for disease development and before disease symptoms are apparent.	For Nassau and Suffolk Counties in New York State, use of this product is limited to 11.5 lb per 1000 sq ft per year (1.95 lb active ingredient per acre).
Downy Rot Leaf Spot Melting-Out	4	14 days	Apply when conditions are favorable for disease development. For downy rot, water in with 3 to 4 gallons of water per 1000 sq ft to increase penetration to crown and roots.	
Dollar Spot	3-8	14-28 days	Apply when conditions are favorable for disease development. Make no more than 3 consecutive applications for dollar spot over before rotating to a registered fungicide with a different mode of action.	
Fusarium Blight	2-8	14-21 days	Apply when conditions are favorable for disease development.	
Fusarium Patch (SBH Snow Mold) Leaf Smut	4-8 4	Fall - Winter 14 days	Apply prior to snow cover. Apply in the fall after grass ceases dormancy and prior to the 1st mow prior to the initiation of growth.	
Neovoss Ring Spot	4-8	Spring: 28 days Fall: 28 days	Make applications on a preventative basis in early to mid-spring. Make 2 applications beginning in August before the turf goes dormant. Apply 6 lb/1000 sq ft followed by a second application one month later.	
Powdery Mildew Rust	4	14-28 days	Apply when conditions are favorable for disease development.	
Summer Patch	4-8	14-28 days	Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending on turf conditions from local turfgrass extension experts. Water in with at least 3 to 4 gallons of water per 1000 sq ft to increase spray penetration to crown and roots.	
Turf Patch	8	Spring/Fall: 28 days	To reduce the severity of turf patch, make 1 to 2 (1) applications in September and October or when night temperatures drop to 50°F and 1 to 2 spring applications in April and May depending on local recommendations.	
Zoysia Large Patch	8	Fall: 28 days	Make applications in fall before turf becomes dormant and water in with at least 3 to 4 gallons of water per 1000 sq ft to increase penetration to crown and roots.	

Application rate is based on commercial turfgrass, or residential turfgrass, or lawn areas of parks, schools, and other sports fields. Golf courses (fairways, rough, tees, greens) and other business and home lawns.

Residential Turfgrass

Disease	LESCO® Eagle® 0.39% Granular Turf Fungicide (8/1000 sq ft)	Application Interval	Instructions	Restrictions
Anthracnose Red Thread Septoria Leaf Spot	4	14-21 days	Apply when conditions are favorable for disease development.	Do not apply more than 46 lb product per 1000 sq ft per year.
Brown Patch		14 days	Begin applications when conditions are favorable for disease development and before disease symptoms are apparent.	For Nassau and Suffolk Counties in New York State, use of this product is limited to 11.5 lb per 1000 sq ft per year (1.95 lb active ingredient per acre).
Downy Rot Zona Leaf Spot		14 days	Apply when conditions are favorable for disease development.	

EPA Reg. No. 62719-461-10404
EPA Est. No. 62757-08-001 (M), 62767-FL-001 (S)
Final letter of label code and other manufacturing info.

#092019

T

Residential Turfgrass continued

Disease	LESCO® Eagle® 0.39% Granular Turf Fungicide (8/1000 sq ft)	Application Interval	Instructions	Restrictions
Downy Rot Leaf Spot Melting-Out		14 days	Apply when conditions are favorable for disease development. For downy rot, water in with 3 to 4 gallons of water per 1000 sq ft to increase penetration to crown and roots.	
Dollar Spot	4	14-21 days	Apply when conditions are favorable for disease development. Make no more than 3 consecutive applications for dollar spot over before rotating to a registered fungicide with a different mode of action.	
Fusarium Blight		14-21 days	Apply when conditions are favorable for disease development.	
Fusarium Patch (SBH Snow Mold) Leaf Smut		Fall - Winter 14 days	Apply prior to snow cover. Apply in the fall after grass enters dormancy and prior to the spring prior to the initiation of growth.	
Neovoss Ring Spot		Spring: 28 days Fall: 28 days	Make applications on a preventative basis in early to mid-spring. Make 2 applications beginning in August before the turf goes dormant.	
Powdery Mildew Rust		14-28 days	Apply when conditions are favorable for disease development.	
Summer Patch		14-28 days	Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending on turf conditions from local turfgrass extension experts. Water in with at least 3 to 4 gallons of water per 1000 sq ft to increase spray penetration to crown and roots.	

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened packages of once to the Seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use, and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Seller MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, lightning, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for disease or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought or used.
2. Replacement of amount of product used.

To the extent permitted by law, Seller shall not be liable for losses or damages resulting from handling or use of this product, unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Information concerning the raw materials comprising this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1385 East 36th Street, Cleveland, Ohio 44114-4114, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.lesco.com/labels/092019>.

Contains Eagle Specialty Fungicide, a registered trademark of Dow. LESCO and Poly Plus are registered trademarks and the revealing design is a trademark of LESCO Technology, LLC. Eagle and Eagle are registered trademarks of Bayer CropScience LLC. Eagle is a registered trademark of C V Van Duyn N.V. Vires is a registered trademark of Rotamex Custom-Plyon Company.

©REGULATORY REVIEW AND LABEL - Winter 2019, 400 (10/2019) Rev: 1/2019 VT

Net Weight: 40 lb (18.18 kg)

F1660

Distributed by: LESCO, Inc.
1385 East 36th Street
Cleveland, OH 44114-4114